



Ubiquitin mouse Monoclonal Antibody(5F1)


Catalog No.	QYA10216A
Size.	100ug
Protein Name.	Ubiquitin
Source.	Mouse
Immunogen.	Synthetic Peptide of Ubiquitin
Purification.	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Specificity.	Ubiquitin protein detects endogenous levels of Ubiquitin
Formulation.	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide..
Concentration.	1 mg/ml
Storage / Stability.	-20 °C/1 year
Human Gene ID.	7314
Human Swiss-Prot No.	PAN
Reactivity.	Human, Rat, Mouse
Applications.	WB, IF,IHC-p
Observed band (KDa)	N/A
Dilution.	WB 1:1000-2000, IHC 1:100-200 IF 1:200




Immunohistochemical analysis of paraffin-embedded Human-Tonsil tissue. 1,Ubiquitin Mouse Monoclonal Antibody(5F1) was diluted at 1:200(4 °C,overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98 °C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.




Immunohistochemical analysis of paraffin-embedded Rat-lung tissue. 1, Ubiquitin Mouse Monoclonal Antibody(5F1) was diluted at 1:200(4 °C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval(>98 °C, 20min). 3, Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.




Immunofluorescence analysis of Human-stomach-cancer tissue. 1, Ubiquitin Mouse Monoclonal Antibody(5F1)(red) was diluted at 1:200(4 °C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of Mouse-brain tissue. 1, Ubiquitin Mouse Monoclonal Antibody(5F1)(red) was diluted at 1:200(4 °C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunofluorescence analysis of Rat-brain tissue. 1, Ubiquitin Mouse Monoclonal Antibody(5F1)(red) was diluted at 1:200(4 °C, overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300(room temperature, 50min). 3, Picture B: DAPI(blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Immunohistochemical analysis of paraffin-embedded Human Breast Carcinoma Tissue using Ubiquitin Mouse mAb diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Human Stomach Carcinoma Tissue using Ubiquitin Mouse mAb diluted at 1:200.